

Rationale Document for EUCAST clinical breakpoints

Agent	Voriconazole	
Current version	4.0	4th February, 2020
Previous versions	3.0	19 th December 2017
	1.0	20 th May 2012 (Voriconazole and <i>Aspergillus</i> spp.)
	2.0	20 th March 2010 (Voriconazole) (vs. <i>Candida</i>)
	1.0	22 nd May 2008 (Voriconazole) (vs. <i>Candida</i>)

Foreword

EUCAST

The European Committee on Antimicrobial Susceptibility Testing (EUCAST) is organised by the European Society for Clinical Microbiology and Infectious Diseases (ESCMID), the European Centre for Disease Prevention and Control (ECDC), and the active national antimicrobial breakpoint committees in Europe. EUCAST was established by ESCMID in 1997, was restructured in 2001-2002 and has been in operation in its current form since 2002. The current remit of EUCAST is to harmonise clinical breakpoints for existing drugs in Europe, to determine clinical breakpoints for new drugs, to set epidemiological (microbiological) breakpoints, to revise breakpoints as required, to harmonise methodology for antimicrobial susceptibility testing, to develop a website with MIC and zone diameter distributions of antimicrobial agents for a wide range of organisms and to liaise with European governmental agencies and European networks involved with antimicrobial resistance and resistance surveillance. Information on EUCAST and EUCAST breakpoints is available on the EUCAST website at <http://www.EUCAST.org>.

EUCAST rationale documents

EUCAST rationale documents summarise the information on which the EUCAST clinical breakpoints are based.

Availability of EUCAST document

All EUCAST documents are freely available from the EUCAST website at <http://www.EUCAST.org>.

Citation of EUCAST documents

The copyright of all documents and data published on the EUCAST website remains with EUCAST. All are freely available for re-use if reference to the EUCAST website is given and documents and data are not resold. Any secondary publication of the data must be referenced with the declaration that "These data have (or this document has) been produced in part under ECDC service contracts, is made available at no cost by EUCAST and can be accessed freely on the EUCAST website www.eucast.org. EUCAST recommendations are frequently updated and the latest versions are available at www.eucast.org."

This rationale document should be cited as: "European Committee on Antimicrobial Susceptibility Testing. Voriconazole: Rationale for the clinical breakpoints, version 4.0, 2020. <http://www.eucast.org>."

1. Introduction

Voriconazole is a triazole antifungal agent with *in vitro* activity against *Candida* spp., *Cryptococcus* spp., *Aspergillus* spp., *Scedosporium apiospermum* and other less common pathogens.

The drug is approved in Europe for the following indications:

- (i) Treatment of invasive aspergillosis.
- (ii) Treatment of candidaemia in non-neutropenic patients.
- (iii) Treatment of fluconazole-resistant serious invasive *Candida* infections (including *C. krusei*).
- (iv) Treatment of serious fungal infections caused by *Scedosporium* spp. and *Fusarium* spp.
- (v) Prophylaxis of invasive fungal infections in high risk allogeneic hematopoietic stem cell transplant (HSCT) recipients.

The mould species most frequently causing human invasive aspergillosis include *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus terreus* and *Aspergillus niger*. The *in vitro* activity of voriconazole against these species of *Aspergillus* is reasonably uniform, but acquired resistance has been reported, even among isolates obtained from triazole naive patients (hence routine susceptibility testing is of utmost importance). *A. fumigatus* is a species complex including rarer sibling species that may exhibit differences in their susceptibility to antifungal agents.

The *Candida* species most frequently involved in causing human infections include *C. albicans*, *C. parapsilosis*, *C. tropicalis*, *C. glabrata* and *C. krusei*. The *in vitro* activity of voriconazole against species of *Candida* is not uniform. MICs are approx. 10 times higher for *C. glabrata* and *C. krusei* than for *C. albicans*. Therefore, every attempt should be made to identify *Candida* to species level. In general, the MICs of voriconazole for fluconazole-resistant isolates are proportionally higher than are those of fluconazole-susceptible isolates.

The European Committee on Antimicrobial Susceptibility Testing Subcommittee on Antifungal Susceptibility Testing (EUCAST-AFST) determined breakpoints for voriconazole against *Aspergillus* spp. and *Candida* spp. in 2012 and 2008, respectively.

In version 4.0 of this rationale document, breakpoints for *Aspergillus nidulans* have been set. Moreover, to encompass with the new definition of the I category, the resistance breakpoints for voriconazole against *Aspergillus fumigatus* has been lowered (one dilution) and an ATU category for *Aspergillus fumigatus* and for *Aspergillus nidulans* have been established.

2. Dosage

		IV (mg/kg/day)	Oral (mg/day unless otherwise stated)
Adults	Loading dose	6 x 2	In >40 kg 400 x 2; In <40 kg 200 x 2
	Maintenance dose	4 x 2	In >40 kg 200 x 2; In <40 kg 100 x 2
Children (2 to <12 years) and young adolescents (12 to 14 years) with low body weight (<50 kg)	Loading dose	9 x 2	NR
	Maintenance dose	8 x 2	9 x 2 (a maximum dose of 350 mg twice daily)
Maximum dose schedule	<p>Adults. If patient response to treatment is inadequate, the oral maintenance dose may be increased to 300 mg twice daily. For patients less than 40 kg the oral dose may be increased to 150 mg twice daily.</p> <p>Children: If patient response to treatment is inadequate, the dose may be increased by 1 mg/kg steps (or by 50 mg steps if the maximum oral dose of 350 mg was used initially).</p>		
Available formulations	Voriconazole is available in the following pharmaceutical forms: film-coated tablets (50 or 200 mg), powder for oral suspension (40 mg/ml), and powder for solution for infusion (200 mg).		
Comments on dosing	<p>The recommended dosing regimen for prophylaxis and treatment are similar.</p> <p>Treatment duration should be as short as possible depending on the patient's clinical and mycological response. Long-term exposure to voriconazole greater than 180 days (6 months) requires careful assessment of the benefit-risk balance. If an adult patient is unable to tolerate treatment at a higher dose, reduce the oral dose by 50 mg steps to the 200 mg twice daily (or 100 mg twice daily for patients less than 40 kg) maintenance dose.</p> <p>On the basis of the high oral bioavailability (96%), switching between intravenous and oral administration is appropriate when clinically indicated. Use in paediatric patients aged 2 to <12 years with hepatic or renal insufficiency has not been studied.</p>		
TDM	The bioavailability of voriconazole is variable due to variable metabolism and potential drug interactions. Dose adjustment based upon therapeutic drug monitoring (TDM) is recommended (trough concentration determination, see below). Voriconazole displays a non-linear kinetics in adults but linear kinetics in children.		

NR = Not recommended

3a. MIC distributions (numbers) and epidemiological cut-off (ECOFF) values (mg/L)

	0.002	0.004	0.008	0.016	0.03	0.06	0.125	0.25	0.5	1	2	4	8	16	32	64	128	256	512	ECOFF
<i>Aspergillus flavus</i>	0	0	0	0	2	4	8	25	46	149	42	4	1	2	0	0	0	0	0	2
<i>Aspergillus fumigatus</i>	0	0	0	0	3	9	85	405	906	288	67	54	44	30	54	0	0	0	0	1
<i>Aspergillus nidulans</i>	0	0	0	3	2	5	18	42	24	6	1	0	1	5	0	0	0	0	0	1
<i>Aspergillus niger</i>	0	0	0	1	1	2	4	13	68	84	35	1	0	1	0	0	0	0	0	2
<i>Aspergillus terreus</i>	0	0	0	0	1	2	2	18	52	154	67	9	1	0	0	0	0	0	0	2
<i>Aspergillus versicolor</i>	0	0	0	0	0	0	2	6	7	7	3	0	1	1	0	0	0	0	0	ND
<i>Aspergillus sydowii</i>	0	0	0	0	1	1	4	11	17	21	0	0	0	0	0	0	0	0	0	ND
<i>Candida albicans</i> *	50	996	5139	5235	1107	530	207	106	63	42	18	13	18	22	78	6	0	0	0	0.03
<i>Candida auris</i> **	0	0	2	0	2	1	17	12	35	37	13	5	0	0	0	0	0	0	2	ND
<i>Candida dubliniensis</i> *	0	0	0	59	38	2	1	0	1	0	0	0	0	0	0	0	0	0	0	0.03
<i>Candida glabrata</i>	0	5	26	115	258	745	1519	1454	855	384	227	196	95	20	2	6	0	0	0	1
<i>Candida guilliermondii</i>	0	0	0	5	11	53	26	16	6	5	3	0	0	0	0	0	0	0	0	ND
<i>Candida krusei</i>	0	0	0	3	3	7	43	179	149	33	6	3	0	1	0	0	0	0	0	1
<i>Candida parapsilosis</i> *	0	6	287	1326	510	197	123	77	23	16	3	2	1	0	0	0	0	0	0	0.06
<i>Candida tropicalis</i> *	0	28	133	602	934	732	274	113	51	32	8	2	10	20	16	3	0	0	0	0.125
<i>Candida kefyr</i>	0	0	5	23	3	2	1	0	0	0	0	0	0	0	0	0	0	0	0	ND
<i>Candida lusitanae</i>	0	0	10	56	14	3	3	4	1	0	0	0	0	0	0	0	0	0	0	ND
<i>Saccharomyces cerevisiae</i>	0	0	0	3	2	12	15	17	8	1	1	0	0	0	0	0	0	0	0	ND
<i>Cryptococcus neoformans</i>	0	0	0	34	44	126	127	101	33	9	3	1	0	1	0	0	0	0	0	0.5
<i>Cryptococcus gattii</i>	0	0	0	1	1	7	11	11	19	8	0	0	0	0	0	0	0	0	0	ND

The table includes EUCAST MIC distributions available at the time breakpoints were set. They represent combined distributions from multiple sources and time periods. The distributions are used to define the epidemiological cut-offs (ECOFF) and give an indication of the MICs for organisms with acquired or mutational resistance mechanisms. They should not be used to infer resistance rates. When there is insufficient evidence, no ECOFFs have been determined (ND). * The majority of the MIC datasets for *C. albicans*, *C. dubliniensis*, *C. parapsilosis* and *C. tropicalis* are truncated at 0.016 mg/L. Data from these truncated datasets also supports the ECOFFs. For *C. albicans* 4149/4237 (98%) MICs are \leq 0.125 mg/L (15 data sets); for *C. parapsilosis* 1325/1362 (97%) MICs are \leq 0.125 mg/L (16 data sets); and for *C. tropicalis* 676/747 (91%) MICs are \leq 0.125 mg/L (14 data sets). The MIC dataset for *Cryptococcus* are truncated at 0.016 mg/L. ** *C. auris* MIC data-set is from a single centre. It includes isolates from several centres in India, but may include genetically related isolates due to the outbreak characteristics of *C. auris* infections. Although not fulfilling the criteria for ECOFF setting, this data is included for general information and comparison given the emergence of this organism worldwide.

3b. MIC distributions (%) and epidemiological cut-off (ECOFF) values (mg/L)

	No.	0.002	0.004	0.008	0.016	0.03	0.06	0.125	0.25	0.5	1	2	4	8	16	32	64	128	256	512	ECOFF
<i>Aspergillus flavus</i>	283	0	0	0	0	1	1	3	9	16	53	15	1	0	1	0	0	0	0	0	2
<i>Aspergillus fumigatus</i>	1945	0	0	0	0	0	0	4	21	47	15	3	3	2	2	3	0	0	0	0	1
<i>Aspergillus nidulans</i>	107	0	0	0	3	2	5	17	39	22	6	1	0	1	5	0	0	0	0	0	1
<i>Aspergillus niger</i>	210	0	0	0	0	0	1	2	6	32	40	17	0	0	0	0	0	0	0	0	2
<i>Aspergillus terreus</i>	306	0	0	0	0	0	1	1	6	17	50	22	3	0	0	0	0	0	0	0	2
<i>Aspergillus versicolor</i>	27	0	0	0	0	0	0	7	22	26	26	11	0	4	4	0	0	0	0	0	ND
<i>Aspergillus sydowii</i>	55	0	0	0	0	2	2	7	20	31	38	0	0	0	0	0	0	0	0	0	ND
<i>Candida albicans</i> *	13630	0	7	38	38	8	4	2	1	0	0	0	0	0	0	1	0	0	0	0	0.03
<i>Candida auris</i> **	126	0	0	2	0	2	1	13	10	28	29	10	4	0	0	0	0	0	0	2	ND
<i>Candida dubliniensis</i> *	101	0	0	0	58	38	2	1	0	1	0	0	0	0	0	0	0	0	0	0	0.03
<i>Candida glabrata</i>	5907	0	0	0	2	4	13	26	25	14	7	4	3	2	0	0	0	0	0	0	1
<i>Candida guilliermondii</i>	125	0	0	0	4	9	42	21	13	5	4	2	0	0	0	0	0	0	0	0	ND
<i>Candida krusei</i>	427	0	0	0	1	1	2	10	42	35	8	1	1	0	0	0	0	0	0	0	1
<i>Candida parapsilosis</i> *	2571	0	0	11	52	20	8	5	3	1	1	0	0	0	0	0	0	0	0	0	0.06
<i>Candida tropicalis</i> *	2958	0	1	4	20	32	25	9	4	2	1	0	0	0	1	1	0	0	0	0	0.125
<i>Candida kefyr</i>	34	0	0	15	68	9	6	3	0	0	0	0	0	0	0	0	0	0	0	0	ND
<i>Candida lusitanae</i>	91	0	0	11	62	15	3	3	4	1	0	0	0	0	0	0	0	0	0	0	ND
<i>Saccharomyces cerevisiae</i>	60	0	0	0	5	3	20	25	28	13	2	2	0	0	0	0	0	0	0	0	ND
<i>Cryptococcus neoformans</i>	478	0	0	0	7	9	26	27	21	7	2	1	0	0	0	0	0	0	0	0	0.5
<i>Cryptococcus gattii</i>	58	0	0	0	2	2	12	19	19	33	14	0	0	0	0	0	0	0	0	0	ND

This table contains a percentage presentation of the MIC data from table 3a. Footnotes for Table 3a apply to this table as well.

4. Breakpoints prior to harmonisation (mg/L) S≤/R>			
	European breakpoints	CLSI ¹	
General breakpoints:			
	NA		
Species related breakpoints:			
	NA	<i>C. albicans</i> <i>C. krusei</i> <i>C. parapsilosis</i> <i>C. tropicalis</i>	S≤0.125 / R>0.5 S≤0.5 / R>1 S≤0.125 / R>0.5 S≤0.125 / R>0.5

NA = Not available

¹CLSI breakpoints converted to European terminology

5. Pharmacokinetics

	Intravenous formulation			Oral formulation		
Dosage	6 mg/kg x 2 on day 1; maintenance dose 4 mg/kg x 2 Steady state in patients with venous haemofiltration (1)	6 mg/kg x 2 on day 1; maintenance dose 4 mg/kg x 2 454 patients with IA (2)	6 mg/kg x 2 on day 1; maintenance dose 3 mg/kg x 2 Healthy volunteers (3, 4)	400 mg x 2 on day 1; maintenance dose 200 mg x 2 Healthy volunteers (3, 4)	400 mg x 2 on day 1; maintenance dose 200 mg x 2 31 AML patients (5)	6 mg/kg x 2 on day 1; 4 mg/kg x 2 days 2-7; maintenance dose 300 mg x 2 454 patients with IA (2)
C _{max} ± SD (mg/L) Mean (CV), [Range]	5.9 ± 2.9		[3-4.7]	[2-2.3]	3.57 ± 1.73	
C _{min} ± SD (mg/L) Mean (CV), [Range]	1.1 ± 0.3	3.10 (52)	[0.39-0.46]	NA	1.40 ± 1.29	1.51 (74)
C _{av} (mg/L) Mean (CV)						
Total body clearance/F ± SD (L/h); Mean (CV)	12.9 ± 6.7		14-35 (F>0.9)*		11.52 ± 8.49	
T _{1/2} ± SD (h), Mean (range)	14.7 ± 6.5		~6		11.31 ± 9.87	
AUC _{24h} ± SD (mg.h/L) total drug; Mean (CV), [Range]	44.8 ± 7.4	100.2 ± 43.08	26	18-22	52.40 ± 37.16	65.8 ± 29.62
Fraction unbound (IQR)	42					
Volume of distribution/F (L/kg)*; Mean (CV), [Range]	2.96 ± 0.55		4.6			
Comments	<p>*Excellent bioavailability in healthy patients (85-92%), but reduced oral bioavailability (60-65%) in some populations, including pediatrics (6, 7). Co-administration with high fat meals decreases absorption (AUC ↓ 35).</p> <ul style="list-style-type: none"> The pharmacokinetics of voriconazole are non-linear in adults likely due to saturation of its metabolism with respect to dose. The variability in plasma concentrations and systemic exposure varies >100-fold between subjects (based on trough concentrations), depending partly on 					

the respective genotype of the hepatic cytochrome P450. CYP2C19 exhibits genetic polymorphism resulting in an approximately 4-fold higher exposure for poor metabolisers than for extensive metabolisers.

- The coefficient of variation of the AUC has been estimated to be 74-100. Comparable exposures are obtained in children with 8 mg/kg x 2 iv or 350 mg x 2 oral ([voriconazole product information](#)). Similar dose adjustments may be required for other patient populations. Protein binding has been estimated to 58% ([voriconazole product information](#)). In a recent study it was however lower in ICU patients 49.6 (42.5-52.5) (8)

6. Pharmacodynamics				
	<i>Aspergillus</i> spp	<i>Candida</i> spp	<i>Cryptococcus</i> spp	
Total drug tAUC ₂₄ :MIC ratio associated with near maximum effect in a dynamic <i>in vitro</i> model of invasive pulmonary aspergillosis	<i>A. fumigatus</i> =32.1 (9)			
fAUC ₂₄ /MIC (95% CI) associated with 50% of maximal effect in an <i>in vitro</i> pharmacokinetic/pharmacodynamic model	<i>A. fumigatus</i> =25.28 (18-35.6) (10)	<i>C. krusei</i> =31 (18-52) <i>C. glabrata</i> =23 (11-47) (11)		
fAUC ₂₄ /MIC±SD (95% CI) associated with 50% survival in non-neutropenic animal model of disseminated aspergillosis	<i>A. fumigatus</i> = 5.86 ^b (12) <i>A. flavus</i> = 14 ± 9 ^c (13)			
fAUC ₂₄ /MIC±SD [range] for achieving ED50		<i>C. albicans</i> = 58.9 ± 66.4 [2-190] (14)		
fAUC/MIC for 2 log reduction in 24 h				
fAUC/MIC from clinical data (CART)				
Trough (C _{min}) concentration associated with an approximately 70% response rate in adult patients	>1-2 mg/L			
Comments	<p>^a fAUC₂₄/CLSI MIC for achieving ED50 is the mean fAUC₂₄/CLSI MIC equivalent to ED50 in mg/kg/24h. ED50 is the dose required to achieve 50% of maximum effect (E_{max}) after 24 h of treatment. The fAUC₂₄/EUCAST MIC was calculated based on the EUCAST MICs of the <i>Candida</i> isolates used by Andes et al (14). The EUCAST MICs were 0.4 (1.6-0) twofold dilutions lower than CLSI MICs.</p> <p>^b the total AUC/EUCAST MIC associated with 50% survival was 17.6 (15). Since the unbound fraction of voriconazole in animals is 29.87% (12), the fAUC/EUCAST MIC is estimated to be 5.86.</p> <p>^c This is the average fAUC/MIC among four <i>A. flavus</i> isolates with voriconazole MICs of 0.25-4 mg/L. A relatively higher exposure was required for the susceptible <i>A. flavus</i> strains in order to produce the same effect compared to <i>A. fumigatus</i>.</p> <ul style="list-style-type: none"> • A trough concentration of >1 mg/L has been associated with a higher probability of a successful outcome in a cohort of patients most of whom had invasive aspergillosis (16) and in the paediatric setting (17). • A trough concentration:MIC ratio of 2-5 (CLSI MIC method) is associated with a higher probability of a successful clinical outcome, but clinical outcomes from both <i>Candida</i> and <i>Aspergillus</i> were considered to derive this endpoint (18). A trough/EUCAST MIC ratio of 1 was associated with 50% of maximal effect in an <i>in vitro</i> pharmacokinetic/pharmacodynamic model of invasive aspergillosis (10). • Pharmacodynamic studies have not been performed for <i>Cryptococcus</i> spp. yet. • Cells are left empty when data are not readily available. 			

7. Monte Carlo simulations and Pk/Pd breakpoints

Aspergillus and *Candida*

The steady state mean \pm SD tAUC_{24s} after a loading dose of 6 mg/kg x 2 on day 1 followed by 1) the i.v. maintenance dose of 4 mg/kg x 2 (100.2 \pm 43.08 mg.h/l), 2) the oral maintenance dose of 300 mg x 2 (65.8 \pm 29.62 m.h/l) or 3) the oral maintenance dose of 200 mg x 2 (52.4 \pm 37.16 mg.h/l) were used in Monte Carlo analysis in order to determine the probability of 5,000 simulated patients attaining the pharmacodynamic targets 60 fAUC₂₄/MIC for *Candida* and 15 fAUC₂₄/MIC for *Aspergillus* considering that the unbound fraction is 58% (2, 5)

MIC (mg/L)	Target attainment for <i>A. fumigatus</i>			Target attainment for <i>C. albicans</i>		
	Oral 200 mg x2	Oral 300 mg x2	iv 4 mg/kg x2	Oral 200 mg x2	Oral 300 mg x2	iv 4 mg/kg x2
0.016	100	100	100	100	100	100
0.03	100	100	100	100	100	100
0.06	100	100	100	99	100	100
0.125	100	100	100	92	100	100
0.25	99	100	100	61	89	99
0.5	92	100	99	21	34	74
1	62	88	74	3	2	15
2	21	34	14	0	0	0
4	3	2	0	0	0	0




Figure 1. Probabilities of target attainment for different dosing regimens of voriconazole against *Candida* and *Aspergillus*.




Figure 2.

Serum trough levels of voriconazole required to attain the pharmacodynamic target for *Candida* ($fAUC_{24}/MIC= 60$) and for *Aspergillus* ($fAUC_{24}/MIC= 15$). The AUC was correlated with trough levels with the following equation $AUC_{12}= 7.011 + 12.687 * \text{trough concentrations}$ (2, 15). The recommended trough level during treatment for aspergillosis is >1-2 mg/L. Isolates with MICs >1 mg/L for *Candida albicans* and >4 mg/L for *Aspergillus* spp would require voriconazole trough levels ≥ 5.5 mg/L which are associated with increased risk of toxicity (19). These align with the selected EUCAST breakpoint for *A. fumigatus* of 1 mg/L (15).

Cryptococcus

Monte Carlo simulations are not available for EUCAST data because there is not a clear Pk/Pd target defined and voriconazole is not approved for the treatment of *Cryptococcus* infections.

8. Clinical data

Aspergillus

Voriconazole is a first-line agent for the treatment of invasive aspergillosis. Use of the currently licensed regimen with at least 1 week of i.v. dosing results in better clinical responses and overall mortality at the end of 12 weeks compared with amphotericin B deoxycholate (20). Voriconazole is the drug of choice for treatment of aspergillosis of the central nervous system (21). Efficacy and safety of voriconazole was compared to voriconazole and anidulafungin combination therapy for invasive aspergillosis in a randomized, double-blind multi-center trial conducted in patients with hematologic malignancies and hematopoietic cell transplantation. Efficacy and safety between both regimens were not statistically significantly different (six week all-cause mortality was 27.5% for voriconazole and 19.3% for combination therapy) (22). In a randomized, double-blind multicenter trial (SECURE trial) for the primary treatment of invasive mould disease, the efficacy and safety of voriconazole was compared to isavuconazole revealing a similar efficacy of both drugs (six week all-cause mortality was 20% with voriconazole) but a higher rate of drug-related events with voriconazole (23).

The frequency of voriconazole resistance is largely undefined, as many centres do not routinely test the susceptibility of their *Aspergillus* isolates. Itraconazole resistance and cross resistance to other triazoles have currently been reported in most European countries, India, Iran, Taiwan, China, Australia, S-America, Africa and the USA (24-30). Most commonly the resistance is linked to point mutations in the target gene *cyp51A* (31-33). However, at some centres a significant proportion of the isolates with elevated voriconazole MICs lack these mutations, suggesting that other mechanisms including upregulated efflux pumps or up-regulation of target production may also play a role (34). Importantly, in some areas, *A. fumigatus* isolates with acquired resistance mechanisms have been increasingly found in the environment and also found in triazole naive patients failing therapy (35, 36). This is probably related to agricultural azole fungicide use, but as *Aspergillus* spores are carried with wind, with compost, plants etc., their presence is not restricted to fungicide treated geographic areas. The voriconazole MICs obtained by EUCAST for isolates with different *cyp51A* point mutations vary according to the underlying mechanism but they are >1 mg/L for the most commonly-identified mutants (alterations at G54, G138, M220, and the environmental phenotypes TR₃₄/L98H and TR₄₆/Y121F/T289A).

Correlation of *in vitro* voriconazole EUCAST MIC data with clinical outcome has been shown in studies comparing outcome between susceptible and molecularly confirmed resistant isolates (37, 38,). If the MIC suggests the isolate may not belong to the wild-type population, voriconazole should not be used.

Candida

Clinical data have been obtained from studies 608 (Global Candidemia Study), 603 (Empirical Therapy Study), 309/604 (Global Rare and Refractory Studies), 301 (Compassionate Use Protocol), and 606 (Emergency Use Protocol-U.S. and Canada).

Voriconazole was administered intravenously with a loading dose of 6 mg/kg every 12 h for the first 24 h, followed by either 3 mg/kg (studies 603 and 608) or 4 mg/kg every 12 h for 3 days, after which patients were given 200 mg oral every 12h. If oral therapy was administered initially, a loading dose of 400 mg every 12 h on day 1 was followed by a maintenance dose of 200 mg twice daily thereafter. The response to voriconazole therapy was determined by the investigator at the end of therapy as either cure, improvement, which were deemed a success or failure. Clinical outcomes at the end of therapy were compared with the MIC of voriconazole for each *Candida* isolated at baseline, i.e. before treatment. MICs were determined using CLSI M27 A2 methodology.

Geometric mean MICs and global response for different *Candida* species were as follows:

Species	No. isolates	Geometric mean MIC (mg/L)	Response
<i>C. albicans</i>	96	0.0164	72%
<i>C. tropicalis</i>	51	0.1283	73%
<i>C. glabrata</i>	47	0.7937	55%
<i>C. parapsilosis</i>	34	0.0266	85%
<i>Candida</i> spp	12	0.0712	92%
<i>C. krusei</i>	9	0.3650	78%

The response was above 72% for infections caused by every species except *C. glabrata* where the percentage response was 55%. Geometric mean MICs were below 0.25 mg/L except for *C. glabrata* and *C. krusei*. CART analysis of response versus MIC or Log₂MIC yielded an MIC value that allowed discrimination between successes and failures. However, the statistical support for this classification tree was limited as the relative error for the best tree was 0.78, the relative risk was 1.09 and the area under ROC curve was 0.6 with a true positive rate of 78% but a false positive rate exceeding 50%.

CART analysis was made separately for infections due to *C. glabrata* because of the lower response rate. CART analysis of outcome versus MIC was unable to produce an interpretable classification tree of MIC values versus successes and failures (39).

In summary, the data at hand support breakpoints that will categorise wild type *C. albicans*, *C. tropicalis* and *C. parapsilosis* susceptible to voriconazole. However, beyond that there is poor statistical support for any clinical correlation between outcome and MIC.

Cryptococcus

There is no clinical data regarding efficacy against *Cryptococcus* infections.

9. *Aspergillus* and *Candida* Clinical breakpoints

Non-species-related breakpoints	<p>Non-species related breakpoints are determined mainly on the basis of Pk/Pd data and are independent of MIC distributions of specific species. They are for use only for species not mentioned in the table or footnotes. In the case of voriconazole, the pharmacokinetics are variable and clinical data for species other than <i>C. albicans</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i> and for isolates with higher MICs are sparse.</p> <p>EUCAST has therefore refrained from determining non-species related breakpoints for voriconazole. There is insufficient evidence to set non-species-related breakpoints against <i>Aspergillus</i> spp.</p>				
Species-related breakpoints	Organism group	MIC breakpoints (mg/L)			Notes
		S ≤	R >	ATU	
	<i>A. fumigatus</i>	1*	1	2	<p>*The susceptibility breakpoint is set provided adequate drug exposure (>1 mg/L) has been confirmed using TDM (Fig. 2 above). An isolate with an MIC = 2 mg/L should be reported as R with the following comment: "In some clinical situations (non-invasive infections forms) voriconazole can be used provided sufficient exposure is ensured" and refer to reference laboratory for CYP51A sequencing and confirmation of MICs.</p>
	<i>A. nidulans</i>	1*	1	2	
	<i>C. albicans</i>	0.06	0.25		<p>A clinical response of 76% was achieved in infections caused by the species listed below when MICs were lower than or equal to the epidemiological cut-offs. Therefore, wild type populations of <i>C. albicans</i>, <i>C. dubliniensis</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i> are considered susceptible. For <i>Candida</i> the intermediate category is introduced to acknowledge that the increased exposure obtained by iv dosing is sufficient (potentially confirmed by TDM). There is not enough information available for the response to voriconazole of infections caused by <i>Candida</i> isolates with higher MICs or infections caused by <i>Cryptococcus</i> species.</p>
	<i>C. dubliniensis</i>	0.06	0.25		
	<i>C. parapsilosis</i>	0.125	0.25		
	<i>C. tropicalis</i>	0.125	0.25		
<p>Breakpoints were based on PK data, microbiological data and patient outcomes from clinical trials. Clinical information suggests that the wild-type population of <i>A. fumigatus</i> is susceptible to voriconazole. No clinical studies have so far presented outcome data for a significant number of cases involving the other species. While there is inadequate clinical information on outcome for wild type population of <i>A. flavus</i>, <i>A. nidulans</i>, <i>A. terreus</i>, the MIC distribution for <i>A. nidulans</i> is similar to those obtained for <i>A. fumigatus</i> but those for <i>A. flavus</i> and <i>A. terreus</i> are slightly higher. Moreover, the fAUC/MIC ratio required for success in animal models is higher for <i>A. flavus</i> than for <i>A. fumigatus</i> (Table 6). Whether that translates into differential clinical outcome remains to be seen. However, for this reason EUCAST has abstained from setting breakpoints for these non-<i>A. fumigatus</i> species (9, 16-18, 40, 41).</p> <p>Increasing data suggest that TDM is an important adjunct for the optimal clinical use of voriconazole against aspergillosis. Most accept that a trough concentration of >1-2 mg/L is a reasonable target. This was originally based on the MIC₉₀ of medically important fungal pathogens,</p>					

	<p>but several recent studies suggest that patients who attain this target have better clinical outcomes and survival. Isolates with a higher MIC require proportionally higher drug exposure to achieve the same effect (Figure 2). While there are no clinical data available that have specifically used EUCAST MIC methodology, preclinical models suggest that a trough: CLSI MIC ratio value of >2-5 is a potential target for therapeutic drug monitoring for treatment of non-wild-type isolates (e.g. MIC 2 mg/L). If the MIC is used in this manner to individualise dosing, the MIC should be repeated to ensure robust estimates are obtained.</p> <p><i>Candida</i> strains with MIC values above the S/I breakpoint are rare among the species that are normally susceptible. The identification and antimicrobial susceptibility testing of any such isolate must be repeated and, if the result is confirmed, the isolate should be sent to a reference laboratory. Isolates with an MIC above the current resistant breakpoint should be reported resistant until evidence has accumulated regarding the clinical response of infections due to such isolates.</p>
Species without breakpoints	<p>There is inadequate clinical information on the clinical outcome for patients infected with wild-type isolates of non-<i>fumigatus Aspergillus</i> species. If the voriconazole exposure-response relationships for these organisms are similar to those for <i>A. fumigatus</i>, the breakpoints for <i>A. fumigatus</i> could be applied to these species. Until these data are available, EUCAST has refrained from setting breakpoints for these species.</p> <p>A 21% lower response to voriconazole in invasive candidiasis caused by <i>C. glabrata</i> has been shown in clinical studies when compared to the response in infections caused by <i>C. albicans</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i>. CART analysis of outcome versus MIC did not find higher MICs to be the variable causing this reduced response so higher MICs were not the explanation for the lower response. Consequently, there is currently insufficient evidence to set clinical breakpoints for <i>C. glabrata</i>.</p> <p>The apparent clinical response in infections caused by <i>C. krusei</i> is similar to that in infections caused by <i>C. albicans</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i>. However, as there were only 9 cases available for analysis, there is currently insufficient evidence to set clinical breakpoints for <i>C. krusei</i>.</p>
Clinical qualifications	<p>The EUCAST AFST considers voriconazole appropriate therapy for the following <i>Aspergillus</i> infections when caused by wild-type <i>A. fumigatus</i> isolates:</p> <ul style="list-style-type: none"> • Primary treatment of invasive aspergillosis including <i>Aspergillus</i> infections of the central nervous system • Treatment of chronic pulmonary aspergillosis <p>The EUCAST AFST considers voriconazole appropriate therapy for the following <i>Candida</i> infections when caused by wild type <i>C. albicans</i>, <i>C. dubliniensis</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i>:</p> <ul style="list-style-type: none"> • Candidaemia in non-neutropenic patients • Invasive candidiasis • Oesophageal candidiasis

Dosage	The EUCAST breakpoints apply to licensed dosing of voriconazole. Dosage escalation may be considered if clinically indicated, and may be guided by therapeutic drug monitoring.
Additional comment	<p>The pharmacokinetics of voriconazole are highly variable. It has been shown that there is a decrease in the response of patients with invasive fungal infections when voriconazole trough blood levels are below 1 mg/L (16, 17, 42). Conversely, voriconazole trough blood levels above 5.5 mg/L have been associated with an increase in toxicity (16). Others report a favourable response when random voriconazole blood concentrations exceed 2.05 mg/L (43). Voriconazole therapeutic drug monitoring is highly recommended to ensure optimal drug exposure. Voriconazole exhibits (pseudo) linear pharmacokinetics in younger children and transitions in adolescence to classical non-linear pharmacokinetics. The probability of elevated liver function tests and central nervous system toxicity is higher with higher drug exposures.</p> <p>Of note, most experience rests with <i>Aspergillus</i> infections and none with invasive cryptococcosis.</p> <p>The EUCAST AFST will review breakpoints for voriconazole when there are more data available for species which were not assigned breakpoints during the present review, when there are clinical data for <i>Candida</i> isolates with MIC values outside the wild-type distribution or when there are data to more closely define therapeutic and toxic levels of voriconazole.</p>

10. Exceptions noted for individual national committees

None.

References

1. Fuhrmann V, Schenk P, Jaeger W, Miksits M, Kneidinger N, Warszawska J, et al. Pharmacokinetics of voriconazole during continuous venovenous haemodiafiltration. *J Antimicrob Chemother.* 2007 Nov;60(5):1085-90.
2. Liu P, Mould DR. Population pharmacokinetic analysis of voriconazole and anidulafungin in adult patients with invasive aspergillosis. *Antimicrob Agents Chemother.* 2014 Aug;58(8):4718-26.
3. Theuretzbacher U, Ihle F, Derendorf H. Pharmacokinetic/pharmacodynamic profile of voriconazole. *Clin Pharmacokinet.* 2006;45(7):649-63.
4. Purkins L, Wood N, Greenhalgh K, Eve MD, Oliver SD, Nichols D. The pharmacokinetics and safety of intravenous voriconazole - a novel wide-spectrum antifungal agent. *Br J Clin Pharmacol.* 2003 Dec;56 Suppl 1:2-9.
5. Geist MJ, Egerer G, Burhenne J, Riedel KD, Weiss J, Mikus G. Steady-state pharmacokinetics and metabolism of voriconazole in patients. *J Antimicrob Chemother.* 2013 Nov;68(11):2592-9.
6. Pascual A, Csajka C, Buclin T, Bolay S, Bille J, Calandra T, et al. Challenging recommended oral and intravenous voriconazole doses for improved efficacy and safety: population pharmacokinetics-based analysis of adult patients with invasive fungal infections. *Clin Infect Dis.* 2012 Aug;55(3):381-90.
7. Scholz I, Oberwittler H, Riedel KD, Burhenne J, Weiss J, Haefeli WE, et al. Pharmacokinetics, metabolism and bioavailability of the triazole antifungal agent voriconazole in relation to CYP2C19 genotype. *Br J Clin Pharmacol.* 2009 Dec;68(6):906-15.
8. Vanstraelen K, Wauters J, De Loor H, Vercammen I, Annaert P, Lagrou K, et al. Protein-binding characteristics of voriconazole determined by high-throughput equilibrium dialysis. *J Pharm Sci.* 2014 Aug;103(8):2565-70.
9. Jeans AR, Howard SJ, Al-Nakeeb Z, Goodwin J, Gregson L, Majithiya JB, et al. Pharmacodynamics of voriconazole in a dynamic in vitro model of invasive pulmonary aspergillosis: implications for in vitro susceptibility breakpoints. *J Infect Dis.* 2012 Aug 01;206(3):442-52.
10. Siopi M, Mavridou E, Mouton JW, Verweij PE, Zerva L, Meletiadis J. Susceptibility breakpoints and target values for therapeutic drug monitoring of voriconazole and *Aspergillus fumigatus* in an in vitro pharmacokinetic/pharmacodynamic model. *J Antimicrob Chemother.* 2014 Jun;69(6):1611-9.
11. Georgiou C, Siopi M, Kanioura L, Zerva L, Arendrup M, Mouton J, et al. Determination of EUCAST PK/PD breakpoints for voriconazole and *C. glabrata*. 26th European Congress on Clinical Microbiology and Infectious Diseases (ECCMID; Abstract no. 5233), Amsterdam, The Netherlands. 2016.
12. Mavridou E, Bruggemann RJ, Melchers WJ, Verweij PE, Mouton JW. Impact of *cyp51A* mutations on the pharmacokinetic and pharmacodynamic properties of voriconazole in a murine model of disseminated aspergillosis. *Antimicrob Agents Chemother.* 2010 Nov;54(11):4758-64.
13. Rudramurthy SM, Seyedmousavi S, Dhaliwal M, Chakrabarti A, Meis JF, Mouton JW. Pharmacodynamics of Voriconazole against Wild-Type and Azole-Resistant *Aspergillus flavus* Isolates in a Nonneutropenic Murine Model of Disseminated Aspergillosis. *Antimicrob Agents Chemother.* 2017 Jan;61(1).
14. Andes D, Marchillo K, Stamstad T, Conklin R. In vivo pharmacokinetics and pharmacodynamics of a new triazole, voriconazole, in a murine candidiasis model. *Antimicrob Agents Chemother.* 2003 Oct;47(10):3165-9.
15. Seyedmousavi S, Mouton JW, Melchers WJ, Bruggemann RJ, Verweij PE. The role of azoles in the management of azole-resistant aspergillosis: from the bench to the bedside. *Drug Resist Updat.* 2014 Jul;17(3):37-50.

16. Pascual A, Calandra T, Bolay S, Buclin T, Bille J, Marchetti O. Voriconazole therapeutic drug monitoring in patients with invasive mycoses improves efficacy and safety outcomes. *Clin Infect Dis*. 2008 Jan 15;46(2):201-11.
17. Neely M, Rushing T, Kovacs A, Jelliffe R, Hoffman J. Voriconazole pharmacokinetics and pharmacodynamics in children. *Clin Infect Dis*. 2010 Jan 01;50(1):27-36.
18. Troke PF, Hockey HP, Hope WW. Observational study of the clinical efficacy of voriconazole and its relationship to plasma concentrations in patients. *Antimicrob Agents Chemother*. 2011 Oct;55(10):4782-8.
19. Lecefel C, Eloy P, Chauvin B, Wyplosz B, Amilien V, Massias L, et al. Worsening pneumonitis due to a pharmacokinetic drug-drug interaction between everolimus and voriconazole in a renal transplant patient. *J Clin Pharm Ther*. 2015 Feb;40(1):119-20.
20. Herbrecht R, Denning DW, Patterson TF, Bennett JE, Greene RE, Oestmann JW, et al. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. *N Engl J Med*. 2002 Aug 08;347(6):408-15.
21. Schwartz S, Ruhnke M, Ribaud P, Corey L, Driscoll T, Cornely OA, et al. Improved outcome in central nervous system aspergillosis, using voriconazole treatment. *Blood*. 2005 Oct 15;106(8):2641-5.
22. Marr KA, Schlamm HT, Herbrecht R, Rottinghaus ST, Bow EJ, Cornely OA, et al. Combination antifungal therapy for invasive aspergillosis: a randomized trial. *Ann Intern Med*. 2015 Jan 20;162(2):81-9.
23. Maertens JA, Raad, II, Marr KA, Patterson TF, Kontoyiannis DP, Cornely OA, et al. Isavuconazole versus voriconazole for primary treatment of invasive mould disease caused by *Aspergillus* and other filamentous fungi (SECURE): a phase 3, randomised-controlled, non-inferiority trial. *Lancet*. 2016 Feb 20;387(10020):760-9.
24. Howard SJ, Cerar D, Anderson MJ, Albarrag A, Fisher MC, Pasqualotto AC, et al. Frequency and evolution of Azole resistance in *Aspergillus fumigatus* associated with treatment failure. *Emerg Infect Dis*. 2009 Jul;15(7):1068-76.
25. Snelders E, van der Lee HA, Kuijpers J, Rijs AJ, Varga J, Samson RA, et al. Emergence of azole resistance in *Aspergillus fumigatus* and spread of a single resistance mechanism. *PLoS Med*. 2008 Nov 11;5(11):e219.
26. Chowdhary A, Kathuria S, Xu J, Sharma C, Sundar G, Singh PK, et al. Clonal expansion and emergence of environmental multiple-triazole-resistant *Aspergillus fumigatus* strains carrying the TR34/L98H mutations in the *cyp51A* gene in India. *PLoS One*. 2012;7(12):e52871.
27. Seyedmousavi S, Hashemi SJ, Zibafar E, Zoll J, Hedayati MT, Mouton JW, et al. Azole-resistant *Aspergillus fumigatus*, Iran. *Emerging infectious diseases*. 2013 May;19(5):832-4.
28. Liu M, Zeng R, Zhang L, Li D, Lv G, Shen Y, et al. Multiple *cyp51A*-based mechanisms identified in azole-resistant isolates of *Aspergillus fumigatus* from China. *Antimicrobial agents and chemotherapy*. 2015 Jul;59(7):4321-5.
29. Wu CJ, Wang HC, Lee JC, Lo HJ, Dai CT, Chou PH, et al. Azole-resistant *Aspergillus fumigatus* isolates carrying TR34/L98H mutations in Taiwan. *Mycoses*. 2015 Sep;58(9):544-9.
30. Wiederhold NP, Gil VG, Gutierrez F, Lindner JR, Albatineh MT, McCarthy DI, et al. First Detection of TR34 L98H and TR46 Y121F T289A *Cyp51* Mutations in *Aspergillus fumigatus* Isolates in the United States. *J Clin Microbiol*. 2016 Jan;54(1):168-71.
31. Mellado E, Garcia-Effron G, Alcazar-Fuoli L, Cuenca-Estrella M, Rodriguez-Tudela JL. Substitutions at methionine 220 in the 14 α -sterol demethylase (*Cyp51A*) of *Aspergillus fumigatus* are responsible for resistance in vitro to azole antifungal drugs. *Antimicrob Agents Chemother*. 2004 Jul;48(7):2747-50.
32. Mellado E, Garcia-Effron G, Alcazar-Fuoli L, Melchers WJ, Verweij PE, Cuenca-Estrella M, et al. A new *Aspergillus fumigatus* resistance mechanism conferring in vitro cross-resistance to azole antifungals involves a combination of *cyp51A* alterations. *Antimicrob Agents Chemother*. 2007 Jun;51(6):1897-904.
33. Mellado E, Garcia-Effron G, Buitrago MJ, Alcazar-Fuoli L, Cuenca-Estrella M, Rodriguez-Tudela JL. Targeted gene disruption of the 14- α sterol demethylase (*cyp51A*) in *Aspergillus fumigatus* and its role in azole drug susceptibility. *Antimicrob Agents Chemother*. 2005 Jun;49(6):2536-8.
34. Arendrup MC, Mavridou E, Mortensen KL, Snelders E, Frimodt-Moller N, Khan H, et al. Development of azole resistance in *Aspergillus fumigatus* during azole therapy associated with change in virulence. *PLoS One*. 2010;5(4):e10080.
35. Snelders E, Huis In 't Veld RA, Rijs AJ, Kema GH, Melchers WJ, Verweij PE. Possible environmental origin of resistance of *Aspergillus fumigatus* to medical triazoles. *Appl Environ Microbiol*. 2009 Jun;75(12):4053-7.
36. Mortensen KL, Mellado E, Lass-Flörl C, Rodriguez-Tudela JL, Johansen HK, Arendrup MC. Environmental study of azole-resistant *Aspergillus fumigatus* and other aspergilli in Austria, Denmark, and Spain. *Antimicrob Agents Chemother*. 2010 Nov;54(11):4545-9.

37. Lestrade PP, Bentvelsen RG, Schauwvlieghe AFAD, Schalekamp S, van der Velden WJFM, Kuiper EJ, van Paassen J, van der Hoven B, van der Lee HA, Melchers WJG, de Haan AF, van der Hoeven HL, Rijnders BJA, van der Beek MT, Verweij PE. Voriconazole Resistance and Mortality in Invasive Aspergillosis: A Multicenter Retrospective Cohort Study. *Clin Infect Dis*. 2019 Apr 24;68(9):1463-1471. doi: 10.1093/cid/ciy859.
38. Resendiz-Sharpe A, Mercier T, Lestrade PPA, van der Beek MT, von dem Borne PA, Cornelissen JJ, De Kort E, Rijnders BJA, Schauwvlieghe AFAD, Verweij PE, Maertens J, Lagrou K. Prevalence of voriconazole-resistant invasive aspergillosis and its impact on mortality in haematology patients. *J Antimicrob Chemother* 2019; 74: 2759–2766.
39. Pfaller MA, Diekema DJ, Rex JH, Espinel-Ingroff A, Johnson EM, Andes D, et al. Correlation of MIC with outcome for *Candida* species tested against voriconazole: analysis and proposal for interpretive breakpoints. *Journal of clinical microbiology*. 2006 Mar;44(3):819-26.
40. Andes D, Pascual A, Marchetti O. Antifungal therapeutic drug monitoring: established and emerging indications. *Antimicrob Agents Chemother*. 2009 Jan;53(1):24-34.
41. Hope WW, Billaud EM, Lestner J, Denning DW. Therapeutic drug monitoring for triazoles. *Curr Opin Infect Dis*. 2008 Dec;21(6):580-6.
42. Dolton MJ, Ray JE, Chen SC, Ng K, Pont LG, McLachlan AJ. Multicenter study of voriconazole pharmacokinetics and therapeutic drug monitoring. *Antimicrob Agents Chemother*. 2012 Sep;56(9):4793-9.
43. Smith J, Safdar N, Knasinski V, Simmons W, Bhavnani SM, Ambrose PG, et al. Voriconazole therapeutic drug monitoring. *Antimicrob Agents Chemother*. 2006 Apr;50(4):1570-2.
- 44.